



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\frac{dy}{d\phi}(R+r)\cos\phi - (R+r)\cos\frac{R+r}{r}\phi.$$

$$\text{But } \frac{ds^2}{d\phi^2} = \frac{dx^2}{d\phi} + \frac{dy^2}{d\phi^2} = 4(R+r)^2 \sin^2 \frac{R}{2r} \phi.$$

\therefore Length of curve between cusp and cusp

$$= 2 \int_0^{2\pi + r/R} (R+r) \sin \frac{R}{2r} \phi = \frac{8r}{R} (R+r).$$

In the case of the hypocycloid we have in its equation only to put $-\tau$ in lieu of r , and by proceeding in the same way we obtain the length of the curve

$$= -\frac{8r}{R} (R-r).$$

AVERAGE AND PROBABILITY.

86. Proposed by L. C. WALKER, Assistant in Mathematics in Leland Stanford, Jr., University, Palo Alto, Cal.


Two points are taken at random in a circular annulus formed by two concentric circles. Find the chance that the straight line joining the points will not cut the inner variable circle.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

Let P, Q be the two random points, MN the chord through P, Q

Let $AC=r$. $CE=w$, $MQ=x$, $PQ=y$, $\angle ACD' = \theta$. CG , the radius of the variable circle $=u$, p =required chance.

An element of the circle at Q is wdw , at P $yd\theta dy$. The limits of u are 0 and r ; of w , r and u ; of x , $2\sqrt{(r^2-w^2)}$ and 0; of y , 0 and x and doubled; of θ , 0 and 2π .



$$\therefore p = \frac{\frac{2}{\pi^2 r^4} \int_0^r \int_u^r \int_0^{2\sqrt{(r^2-w^2)}} \int_0^x \int_0^{2\pi} du dw dx dy d\theta}{\int_0^r du}$$

$$= \frac{4}{\pi r^5} \int_0^r \int_u^r \int_0^{2\sqrt{(r^2-w^2)}} \int_0^x du dw dx dy$$

$$\begin{aligned}
 &= \frac{2}{\pi r^5} \int_0^r \int_u^r \int_0^{\sqrt{(r^2-w^2)}} x^2 du dw dx \\
 &= \frac{16}{3\pi r^5} \int_0^r \int_u^r (r^2-w^2)^{\frac{3}{2}} du dw \\
 &= \frac{1}{3\pi r^5} \int_0^r [3\pi r^4 - 10r^2 u \sqrt{(r^2-u^2)} + 4u^3 \sqrt{(r^2-u^2)} - 6r^4 \sin^{-1} \frac{u}{r}] du \\
 &= \frac{16}{15\pi} = .3395.
 \end{aligned}$$

87. Proposed by G. B. M. ZERE, A. M., Ph. D., Professor of Science and Mathematics, Chester High School, Chester, Pa.

Find the mean distance of a random point in a sphere from a point, (1) within, (2) without the sphere.

Solution by the PROPOSER.

Let B be the given point. $AB=c$, $AC=a$, $AD=\pm x$.


Then $DB=c-x=BC$, $CE=\sqrt{(a^2-AE^2)}$
 $=\sqrt{[(c-x)^2-(c-AE)^2]}.$

$$\therefore AE = \frac{(a^2+2cx-x^2)}{2c},$$

$$DE=AE-x=\frac{(a^2-x^2)}{2c}.$$

Area of film $CDM=2\pi DB.DE=2\pi(c-x)(a^2-x^2)/2c$.

Also let $BG=y$. Then the area of the surface of this sphere is $4\pi y^2$.



$$(1). \text{ Distance } D = \frac{\int_{-a}^{2c-a} 2\pi DB^2.DE dx + \int_0^{a-c} 4\pi y^3 dy}{\frac{4}{3}\pi a^3}$$

$$\therefore D = \frac{\frac{3}{4a^3c} \int_{-a}^{2c-a} (c-x)^2(a^2-x^2) dx + \frac{3}{a^3} \int_0^{a-c} y^3 dy}{\frac{4}{3}\pi a^3}.$$

$$\therefore D = \frac{3}{4}a + \frac{c^2}{2a} - \frac{c^4}{20a^3}.$$

$$(2). D_1 = \frac{\int_{-a}^a 2\pi DB^2.DE dx}{\frac{4}{3}\pi a^3} = \frac{3}{4a^3c} \int_{-a}^a (c-x)^2(a^2-x^2) dx.$$

$$\therefore D_1 = c + (a^2/5c).$$